




OFFICIAL PUBLICATION OF
THE MINISTRY OF HEALTH,
BRUNEI DARUSSALAM

Brunei International Medical Journal

Volume 14

29 May 2018 (123 Ramadhan 1439H)



Brunei Int Med J. 2018;14:73-76

Brunei International Medical Journal (BIMJ)

Official Publication of the Ministry of Health, Brunei Darussalam

EDITORIAL BOARD

Editor-in-Chief	William Chee Fui CHONG
Sub-Editors	Vui Heng CHONG Ketan PANDE
Editorial Board Members	Nazar LUQMAN Muhd Syafiq ABDULLAH Alice Moi Ling YONG Ahmad Yazid ABDUL WAHAB Jackson Chee Seng TAN Dipo OLABUMUYI Pemasiri Upali TELISINGHE Roselina YAAKUB Pengiran Khairol Asmee PENGIRAN SABTU Dayangku Siti Nur Ashikin PENGIRAN TENGAH

INTERNATIONAL EDITORIAL BOARD MEMBERS

Lawrence HO Khok Yu (Singapore)	Surinderpal S BIRRING (United Kingdom)
Emily Felicia Jan Ee SHEN (Singapore)	Leslie GOH (United Kingdom)
John YAP (United Kingdom)	Chuen Neng LEE (Singapore)
Christopher HAYWARD (Australia)	Jimmy SO (Singapore)
Jose F LAPENA (Philippines)	Simon Peter FROSTICK (United Kingdom)

Advisor

Wilfred PEH (Singapore)

Past Editors

Nagamuttu RAVINDRANATHAN
Kenneth Yuh Yen KOK

Proof reader

John WOLSTENHOLME (CfBT Brunei Darussalam)

Aim and Scope of Brunei International Medical Journal

The Brunei International Medical Journal (BIMJ) is a six monthly peer reviewed official publication of the Ministry of Health under the auspices of the Clinical Research Unit, Ministry of Health, Brunei Darussalam.

The BIMJ publishes articles ranging from original research papers, review articles, medical practice papers, special reports, audits, case reports, images of interest, education and technical/innovation papers, editorials, commentaries and letters to the Editor. Topics of interest include all subjects that relate to clinical practice and research in all branches of medicine, basic and clinical including topics related to allied health care fields. The BIMJ welcomes manuscripts from contributors, but usually solicits reviews articles and special reports. Proposals for review papers can be sent to the Managing Editor directly. Please refer to the contact information of the Editorial Office.

Instruction to authors

Manuscript submissions

All manuscripts should be sent to the Managing Editor, BIMJ, Ministry of Health, Brunei Darussalam; e-mail: editor-in-chief@bimjonline.com. Subsequent correspondence between the BIMJ and authors will, as far as possible via should be conducted via email quoting the reference number.

Conditions

Submission of an article for consideration for publication implies the transfer of the copyright from the authors to the BIMJ upon acceptance. The final decision of acceptance rests with the Editor-in-Chief. All accepted papers become the permanent property of the BIMJ and may not be published elsewhere without written permission from the BIMJ.

Ethics

Ethical considerations will be taken into account in the assessment of papers that have experimental investigations of human or animal subjects. Authors should state clearly in the Materials and Methods section of the manuscript that institutional review board has approved the project. Those investigators without such review boards should ensure that the principles outlined in the Declaration of Helsinki have been followed.

Manuscript categories

Original articles

These include controlled trials, interventional studies, studies of screening and diagnostic tests, outcome studies, cost-effectiveness analyses, and large-scale epidemiological studies. Manuscript should include the following; introduction, materials and methods, results and conclusion. The objective should be stated clearly in the introduction. The text should not exceed 2500 words and references not more than 30.

Review articles

These are, in general, invited papers, but unsolicited reviews, if of good quality, may be considered. Reviews are systematic critical assessments of

literature and data sources pertaining to clinical topics, emphasising factors such as cause, diagnosis, prognosis, therapy, or prevention. Reviews should be made relevant to our local setting and preferably supported by local data. The text should not exceed 3000 words and references not more than 40.

Special Reports

This section usually consist of invited reports that have significant impact on healthcare practice and usually cover disease outbreaks, management guidelines or policy statement paper.

Audits

Audits of relevant topics generally follow the same format as original article and the text should not exceed 1,500 words and references not more than 20.

Case reports

Case reports should highlight interesting rare cases or provide good learning points. The text should not exceed 1000 words; the number of tables, figures, or both should not be more than two, and references should not be more than 15.

Education section

This section includes papers (i.e. how to interpret ECG or chest radiography) with particular aim of broadening knowledge or serve as revision materials. Papers will usually be invited but well written paper on relevant topics may be accepted. The text should not exceed 1500 words and should include not more than 15 figures illustration and references should not be more than 15.

Images of interest

These are papers presenting unique clinical encounters that are illustrated by photographs, radiographs, or other figures. Image of interest should include a brief description of the case and discussion with educational aspects. Alternatively, a mini quiz can be presented and answers will be posted in a different section of the publication. A maximum of

three relevant references should be included. Only images of high quality (at least 300dpi) will be acceptable.

Technical innovations

This section include papers looking at novel or new techniques that have been developed or introduced to the local setting. The text should not exceed 1000 words and should include not more than 10 figures illustration and references should not be more than 10.

Letters to the Editor

Letters discussing a recent article published in the BIMJ are welcome and should be sent to the Editorial Office by e-mail. The text should not exceed 250 words; have no more than one figure or table, and five references.

Criteria for manuscripts

Manuscripts submitted to the BIMJ should meet the following criteria: the content is original; the writing is clear; the study methods are appropriate; the data are valid; the conclusions are reasonable and supported by the data; the information is important; and the topic has general medical interest. Manuscripts will be accepted only if both their contents and style meet the standards required by the BIMJ.

Authorship information

Designate one corresponding author and provide a complete address, telephone and fax numbers, and e-mail address. The number of authors of each paper should not be more than twelve; a greater number requires justification. Authors may add a publishable footnote explaining order of authorship.

Group authorship

If authorship is attributed to a group (either solely or in addition to one or more individual authors), all members of the group must meet the full criteria and requirements for authorship described in the following paragraphs. One or more authors may take responsibility 'for' a group, in which case the other group members are not authors, but may be listed in an acknowledgement.

Authorship requirement

When the BIMJ accepts a paper for publication, authors will be asked to sign statements on (1) financial disclosure, (2) conflict of interest and (3) copyright transfer. The correspondence author may sign on behalf of co-authors.

Authorship criteria and responsibility

All authors must meet the following criteria: to have participated sufficiently in the work to take public responsibility for the content; to have made substantial contributions to the conception and de-

sign, and the analysis and interpretation of the data (where applicable); to have made substantial contributions to the writing or revision of the manuscript; and to have reviewed the final version of the submitted manuscript and approved it for publication. Authors will be asked to certify that their contribution represents valid work and that neither the manuscript nor one with substantially similar content under their authorship has been published or is being considered for publication elsewhere, except as described in an attachment. If requested, authors shall provide the data on which the manuscript is based for examination by the editors or their assignees.

Financial disclosure or conflict of interest

Any affiliation with or involvement in any organisation or entity with a direct financial interest in the subject matter or materials discussed in the manuscript should be disclosed in an attachment. Any financial or material support should be identified in the manuscript.

Copyright transfer

In consideration of the action of the BIMJ in reviewing and editing a submission, the author/s will transfer, assign, or otherwise convey all copyright ownership to the Clinical Research Unit, RIPAS Hospital, Ministry of Health in the event that such work is published by the BIMJ.

Acknowledgements

Only persons who have made substantial contributions but who do not fulfill the authorship criteria should be acknowledged.

Accepted manuscripts

Authors will be informed of acceptances and accepted manuscripts will be sent for copyediting. During copyediting, there may be some changes made to accommodate the style of journal format. Attempts will be made to ensure that the overall meaning of the texts are not altered. Authors will be informed by email of the estimated time of publication. Authors may be requested to provide raw data, especially those presented in graph such as bar charts or figures so that presentations can be constructed following the format and style of the journal. Proofs will be sent to authors to check for any mistakes made during copyediting. Authors are usually given 72 hours to return the proof. No response will be taken as no further corrections required. Corrections should be kept to a minimum. Otherwise, it may cause delay in publication.

Offprint

Contributors will not be given any offprint of their published articles. Contributors can obtain an electronic reprint from the journal website.

DISCLAIMER

All articles published, including editorials and letters, represent the opinion of the contributors and do not reflect the official view or policy of the Clinical Research Unit, the Ministry of Health or the institutions with which the contributors are affiliated to unless this is clearly stated. The appearance of advertisement does not necessarily constitute endorsement by the Clinical Research Unit or Ministry of Health, Brunei Darussalam. Furthermore, the publisher cannot accept responsibility for the correctness or accuracy of the advertisers' text and/or claim or any opinion expressed.

Neglected Osteosarcoma of the Knee in a Young Female Adolescent: A Case Report.

Muhammad Bayu Zohari HUTAGALUNG, Safrizal RAHMAN, Azharuddin AZHARUDDIN
Department of Orthopaedic and Traumatology, Faculty of Medicine, Syiah Kuala University/Dr. Zainoel Abidin Hospital, JL. T Daud Beureueh No 108, Banda Aceh 23124, Indonesia

ABSTRACT

A 16-years old girl presented to the orthopaedic outpatient clinic of Dr. Zainoel Abidin Hospital, Banda Aceh, Indonesia with a progressive growing mass around the left knee since 12 months ago. Diagnostic assesment includes clinical, radiological and fine needle aspiration cytology of the mass confirmed an osteosarcoma. Patients undergo above knee amputation procedure as definitive treatment. Neglected osteosarcoma is a primary malignant bone tumor with more than 3 months delayed from the original time of signs and symptoms occurrence to first presentation for medical care. The combination of unawareness, low socio-economic level and inability to access healthcare facilities are major factors contributing to delayed presentation in this case. Management of neglected osteosarcoma in our region faced a variety of issues such as lack of funding, inadequate diagnostic facilities and limited access to advanced treatment options combined with patient's cultural and socio-economic preferences with regards to management and treatment options.

Keywords: Amputation, Health services accessibility, Osteosarcoma, neglected, Disease management.

INTRODUCTION

Osteosarcoma (OS) is the most common primary malignant bone tumor, usually occurring in the second decade of life.¹ The management of patients with osteosarcoma in developing countries particularly in our geographical region is challenging and often delayed due to the lack of awareness amongst patients of their illness, patients' cultural preferences and poor socio-economic status, coupled with inadequate diagnostic facilities and effective therapy issues.² These factors often led to delayed presentation of malignant bone and soft tissue tumours to

medical care, more recently labeled as neglected malignant bone tumours, when presentation to medical care is delayed by more than 3 months from time of signs and symptoms.^{3,4} Neglected malignant bone tumours because of their advance stage tend to be associated with a poorer prognosis with high morbidity such as limb loss and mortality.³ Clinical presentation such as severe pain, tumor fungation, sepsis, thrombosis, hemorrhage, pathologic fractures and severe functional impairment is often the case. We present here a case of neglected advanced osteosarcoma of the left knee in a young teenage girl and discussed the management options available.

Corresponding author: Muhammad Bayu Zohari Hutagalung, MD, Department of Orthopaedic and Traumatology, Faculty of Medicine, Syiah Kuala University/Dr. Zainoel Abidin Hospital, JL. T Daud Beureueh No 108, Banda Aceh 23124, Indonesia.
Tel: +6282364622502,
email: bayuzohari@gmail.com

CASE REPORT

A 16-years old girl presented to the Orthopae-




Figure 1. A) Mass around the left knee joint; B) Plain x-ray of the mass (Left Knee Joint)

dic Outpatient clinic of Dr. Zainoel Abidin Hospital, Banda Aceh, Indonesia with a progressively enlarging tumour around the left knee of 12 months duration. The patient suffered from severe pain in the knee with inability to walk normally since 7 months ago. She gave a history of previous left lower leg trauma a year ago which was treated by a traditional bone-setter using massage and traditional herbal medicine around the injured knee.

Physical examination showed poor general condition with underweight appearance. Vital signs were within normal limit. There was a large solid tumour measuring 39 cm by 32 cm, arising from lower third of the thigh and extending across the left knee to mid shin. The skin overlying the mass was thin and ulcerated in some areas with underlying dilated veins (Figure 1a). The range of motion of left knee was severely limited.

Blood investigations showed mild anemia with Hb of 9.0 mg/dL and an elevated white cell Count of $18.3 \times 10^3/\text{mm}^3$ and platelets of $646 \times 10^3/\text{U/L}$. Plain radiograph of the left knee showed well-defined soft tissue tumour arising from the upper third of the left tibia with expansile lytic lesions, periosteal reaction (Codman triangle), sunray appearance, lateral shifting of tibia-fibula

bone and extending upwards to the lower third of the femur and downwards to the mid shin level (Figure 1B). Fine needle aspiration cytology of the tumour showed clusters of osteocytes with hyperchromatic pleomorphic nuclei, increased Nuclear-Cytoplasmic (NC) ratio and producing osteoid (Figure 2).

Given the advanced stage of disease, a plain chest radiograph and ultrasonography of abdominal organs were performed for the initial staging of her disease. The chest radiograph was clear and ultrasonography showed no liver or intraabdominal metastatic lesion. She underwent above knee amputation as a definitive treatment for the advance osteosarcoma. She was followed up postoperatively after her discharge up to a year where she was well but had refused to undergo post-operative CT staging or adjuvant chemotherapy.

DISCUSSION

Osteosarcoma is primary mesenchymal malignancy characterized by the production of osteoid.¹ The incidence of Osteosarcoma showed a bimodal peak in life, the first peak occurs in the adolescent and young adult and a second peak in the 6th decade of life.² Even though, there have been significant improvements in the diagnosis and treatment of osteosarcoma,




Figure 2. Fine Needle Aspiration smears of Osteosarcoma (MGG stain, original magnification x25)

overall survival rates are relatively constant for the past 2 decades. The extreme variability of osteosarcoma cases from one to another, making single treatment pathway seems unlikely to be able to treat all or even most of the cases.²

The occurrence of neglected malignant bone tumors such as neglected osteosarcoma in developing countries as in our case is due to a combination of several epidemiological factors such as; unawareness, low socio-economic status and lack of diagnostic facilities and therapeutic options. The lack of awareness of the severity of the illness on the part of the patients combined with patients' cultural or religious beliefs and preferences also contribute to a large extent to the delay in presentation for medical care. The effect of the extremity tumors, especially in lower limbs significantly impaired the mobility and decreased the patient's quality of life.⁵

The most frequent sites of osteosarcoma include the metaphyseal bone of the distal femur, the proximal tibia, and the proximal humerus. The patients with neglected osteosarcoma typically present with enlargement of the tumor with either ulceration of the skin or pain. Initially, pain may be related to activity, but progresses to rest pain or

even night pain which unfortunately is commonly attributed to "growing pains." The median time from onset of symptoms to diagnosis is 4 months, though significant variability exists especially in underdeveloped or developing countries. Mild blunt trauma is often reported as an antecedent event but currently, there is no significant correlation between trauma and osteosarcoma.²

Radiological imaging is the gold standard diagnostic options for assessing extent and stage of malignant bony tumours such as osteosarcoma. Plain radiographs of the involved bone and adjacent joint have become a popular option due to low cost with high diagnostic values. Typical osteosarcoma radiographs appear as a mixed lytic and radiodense lesion adjacent to the metaphyseal bone in an eccentric manner. Cortical destruction and periosteal reaction are common, and typically manifest in a sunburst pattern. There is frequently mass extension into surrounding tissue.² In our institution, Computer Tomography (CT) imaging is not routinely performed and thus tumour staging for distant metastases are based on clinical examination, an ultrasound exam of the abdomen, and plain radiographs of the chest. Fine needle aspiration cytology (FNAC) is a mandatory diagnostic tool in diagnosing bone tumor lesions. The advantages of FNAC for diagnosing bone tumor includes simple, inexpensive, does not require general anesthesia, allows preliminary diagnosis within 15–20 min of aspiration with minimal possibility of seeding the tumor cells. The sensitivity of FNAC for categorizing bone tumors into benign and malignant was 94.7%. For soft tissue tumors, FNAC had a sensitivity of 90.9% and specificity was 100%.⁶ In majority cases of osteosarcoma, presence of osteoid or osteoid-like material together with pleomorphic cells is highly diagnostic of osteosarcoma. In the majority of osteosarcoma especially in the neglected case, the combined investigation of clinical and radiological data together with the FNAC result has been

sufficient for making treatment decisions.⁷

The current standard of care for neglected osteosarcoma is the combination of chemotherapy, complete surgical resection of the primary disease, evaluation of possible metastasis from the primary tumor and its related treatment.^{8,9} Complete surgical resection of the tumor should be the goal for curative treatment in osteosarcoma regardless of the location. In the malignant bone tumor, amputation has been reported as a good oncologic procedure, especially in neglected case. Despite this, amputation is often misconstrued to be a procedure with poor functional results. With the use of excellent prosthesis patients with osteosarcoma who have undergone lower limb amputations showed the ability to sustain a much higher activity level, including impact activities such as running.^{2,8,10}

In the developing countries, treatment options are relatively limited with far less access to the more advanced treatment care compared to a developed healthcare system. The challenge in treating neglected osteosarcoma case in developing countries not just influenced by the a combination of advanced pathology due to delayed treatment and unawareness of the patient, poor access to investigative modalities either financial or lack of diagnostic tools, cultural reluctance to comply with Western treatment related to local culture, limited access to adjuvant therapy and lack of modern surgical limb salvage techniques.¹¹

CONCLUSION

The health-seeking behavior in neglected osteosarcoma patient resulted from the combination of unawareness, low education, low socio-economic level, unable to access healthcare provider and others. An adequate diagnostic and early treatment pathway is needed for patients presenting with neglected

malignant bone tumours in order to improve the quality of life of these patients.

REFERENCES

- 1: Rastogi S, Aggarwal A, Tiwari A, Sharma V. Chemotherapy in Nonmetastatic Osteosarcoma: Recent Advances and Implications for Developing Countries. *J Glob Oncol.* 2017;1–5.
- 2: Geller DS, Gorlick R. Osteosarcoma: A review of diagnosis, management, and treatment strategies. *Clin Adv Hematol Oncol.* 2010;8(10):705–18.
- 3: Discepolo F, Powell TI, Nahal A. Telangiectatic osteosarcoma: Radiologic and pathologic findings. *Radiographics* 2009;29(2):380–3.
- 4: Phatak SV, Kolwadkar PK, Lawange A, Rajderkar D. Telangiectatic osteosarcoma: A case report. *Ind J Radiol Imag* 2005;3:353–4.
- 5: Siddiqui Y, Sherwani M, Khan A, Zahid M, Abbas M, Asif N. Neglected orthopedic oncology - Causes, epidemiology and challenges for management in developing countries. *Indian J Cancer [Internet].* 2015;52(3):325.
- 6: Kaur R, Handa U, Kundu R, Garg SK, Mohan H. Role of fine-needle aspiration cytology and core needle biopsy in diagnosing musculoskeletal neoplasms. *J Cytol.* 2016 Jan-Mar; 33(1): 7–12
- 7: Mohamed Aly A, Shaaban HM, Abou-Sinna I. Accuracy of fine needle aspiration cytology in the diagnosis of bone lesions with radiological assistance: Experience from the National Cancer Institute, Cairo University, Egypt. *Egypt J Radiol Nucl Med.* 2014;45(1):127–35.
- 8: Reed DR, Hayashi M, Wagner L, Binitie O, Stepan DA, Brohl AS, et al. Treatment pathway of bone sarcoma in children, adolescents, and young adults. *Cancer.* 2017;123(12):2206–18.
- 9: Isakoff MS, Bielack SS, Meltzer P, Gorlick R. Osteosarcoma: Current treatment and a collaborative pathway to success. *J Clin Oncol.* 2015;33(27):3029–35.
- 10: Durfee RA, Mohammed M, Luu HH. Review of Osteosarcoma and Current Management. *Rheumatol Ther [Internet].* 2016;3(2):221–43. Available from: <http://link.springer.com/10.1007/s40744-016-0046-y>
- 11: Noor S, Pormósson HS, Zervas CT, Ly T, Gollogly J. Limb versus life - The outcomes of osteosarcoma in Cambodia. *Int Orthop.* 2014;38(3):579–85.