

**(Refer to page 9)**

### **ANSWER: EMPHYSEMATOUS CELLULITIS**

The radiograph (panel) of the left foot shows gas pockets (arrows) located at the 4<sup>th</sup> and 5<sup>th</sup> toes with changes of osteomyelitis affecting the 5<sup>th</sup> toe. Pus and tissue cultures isolated *Morganella morganii* (*M. morganii*) sensitive to the antibiotics (intravenous amoxicillin-clavulanic acid and ceftazidime) given to the patient. The patient was treated with wound debridement disarticulation of the 4<sup>th</sup> and 5<sup>th</sup> toes but later required a below knee amputation to control the infection.



Diabetic foot is a common complication among patients with poorly controlled diabetes mellitus and is the leading cause of limb amputation.<sup>1, 2</sup> Emphysematous cellulitis is uncommon and is potentially catastrophic resulting in amputation and even death. The causative organisms can be *Clostridia* or non-*clostridia* species. Gas formation caused by non-clostridia organisms is due to mixed aerobic (sometimes anaerobic bacteria) metabolism and usually has a more gradual progression and better prognosis compared to *Clostridial* infection.<sup>3</sup> In patient with diabetes mellitus, gas formation can be due to fermentation of organic acid as result of hyperglycemia. Microvascular complications contribute to tissue ischaemia. *M. morganii* is a gram negative bacillus and it can produce severe

local and systemic infection in diabetes patient which can result poor outcomes.<sup>4</sup>

Emphysematous cellulitis requires early diagnosis and should be treated appropriately with glycaemia control and good antibiotic coverage, typically intravenous in the initial phase that can be changed to oral formulation. The antibiotics coverage should be guided by microbial sensitivity results. Surgical interventions (i.e. debridement) may need to be considered early because amputation can greatly effect on the outcome and on the daily life of the patient. In this present case, the infection was not controlled with debridement and disarticulation necessitating a below knee amputation.

### **REFERENCES**

- 1: Cooney DR, Cooney NL. [Gas gangrene and osteomyelitis of the foot in a diabetic patient treated with tea tree oil.](#) *Int J Emerg Med.* 2011; 4:14. [Accessed on 2019 November 1].
- 2: Kuy S, Romero RA, Kuy S. Gas gangrene of the diabetic foot. *J Louisiana State Med Society: official organ of the Louisiana State Medical Society.* 2015; 167:213-4.
- 3: Kono S NR, Arata J, Lipsky BA. [Massive gas-forming gangrene in a diabetic foot infection.](#) *Clin Research Foot Ankle* 2014; 2:i101. [Accessed on 2019 November 1].
- 4: Ghosh S, Bal AM, Malik I, Collier A. [Fatal Morganella morganii bacteraemia in a diabetic patient with gas gangrene.](#) *J Med Microbiol.* 2009; 58:965-7. [Accessed on 2019 November 1].