

Brunei International Medical Journal

OFFICIAL PUBLICATION OF
THE MINISTRY OF HEALTH
AND
UNIVERSITI BRUNEI DARUSSALAM

Volume 18

30 March 2022 (26 Sya'ban 1443H)

KIDNEY HEALTH FOR ALL: BRIDGING THE GAP IN KIDNEY HEALTH EDUCATION AND LITERACY.

Robyn G. LANGHAM¹, Kamyar KALANTAR-ZADEH², Ann BONNER³, Alessandro BALDUCCI⁴, Li-Li HSIAO⁵, Latha A. KUMARASWAMI⁶, Paul LAFFIN⁷, Vassilios LIAKOPOULOS⁸, Gamal SAADI⁹, Eka-mol TANTISATTAMO², Ifeoma ULASI¹⁰ and Siu-Fai LUI¹¹ for the World Kidney Day Joint Steering Committee¹².

ABSTRACT

The high burden of kidney disease, global disparities in kidney care, and poor outcomes of kidney failure bring a concomitant growing burden to persons affected, their families, and carers, and the community at large. Health literacy is the degree to which persons and organizations have or equitably enable individuals to have the ability to find, understand, and use information and services to make informed health-related decisions and actions for themselves and others. Rather than viewing health literacy as a patient deficit, improving health literacy largely rests with health care providers communicating and educating effectively in codesigned partnership with those with kidney disease. For kidney policy makers, health literacy provides the imperative to shift organizations to a culture that places the person at the center of health care. The growing capability of and access to technology provides new opportunities to enhance education and awareness of kidney disease for all stakeholders. Advances in telecommunication, including social media platforms, can be leveraged to enhance persons' and providers' education; The World Kidney Day declares 2022 as the year of "Kidney Health for All" to promote global teamwork in advancing strategies in bridging the gap in kidney health education and literacy. Kidney organizations should work toward shifting the patient-deficit health literacy narrative to that of being the responsibility of health care providers and health policy makers. By engaging in and supporting kidney health-centered policy making, community health planning, and health literacy approaches for all, the kidney communities strive to prevent kidney diseases and enable living well with kidney disease.

KEYWORDS: Educational gap, Empowerment, Health literacy, Health policy, Information technology, Kidney health, Partnership, Prevention, Social media.

Brunei Int Med J. 2022;18:51-62

Brunei International Medical Journal (BIMJ) Official Publication of The Ministry of Health and Universiti Brunei Darussalam

EDITORIAL BOARD

Editor-in-Chief	Ketan PANDE
Sub-Editors	Vui Heng CHONG William Chee Fui CHONG
Editorial Board Members	Muhd Syafiq ABDULLAH Alice Moi Ling YONG Ahmad Yazid ABDUL WAHAB Jackson Chee Seng TAN Pemasiri Upali TELISINGHE Pengiran Khairol Asmee PENGIRAN SABTU Dayangku Siti Nur Ashikin PENGIRAN TENGAH

INTERNATIONAL EDITORIAL BOARD MEMBERS

Lawrence HO Khek Yu (Singapore)	Chuen Neng LEE (Singapore)
Wilfred PEH (Singapore)	Emily Felicia Jan Ee SHEN (Singapore)
Surinderpal S BIRRING (United Kingdom)	Leslie GOH (United Kingdom)
John YAP (United Kingdom)	Ian BICKLE (United Kingdom)
Nazar LUQMAN (Australia)	Christopher HAYWARD (Australia)
Jose F LAPENA (Philippines)	

Advisor

Wilfred PEH (Singapore)

Past Editors-in-Chief

Nagamuttu RAVINDRANATHAN
Kenneth Yuh Yen KOK
Chong Vui Heng
William Chong Chee Fui

Proof reader

John WOLSTENHOLME (CfBT Brunei Darussalam)

Aim and Scope of Brunei International Medical Journal

The Brunei International Medical Journal (BIMJ) is a six monthly peer reviewed official publication of the Ministry of Health under the auspices of the Clinical Research Unit, Ministry of Health, Brunei Darussalam.

The BIMJ publishes articles ranging from original research papers, review articles, medical practice papers, special reports, audits, case reports, images of interest, education and technical/innovation papers, editorials, commentaries and letters to the Editor. Topics of interest include all subjects that relate to clinical practice and research in all branches of medicine, basic and clinical including topics related to allied health care fields. The BIMJ welcomes manuscripts from contributors, but usually solicits reviews articles and special reports. Proposals for review papers can be sent to the Managing Editor directly. Please refer to the contact information of the Editorial Office.

Instruction to authors

Manuscript submissions

All manuscripts should be sent to the Managing Editor, BIMJ, Ministry of Health, Brunei Darussalam; e-mail: editor-in-chief@bimjonline.com. Subsequent correspondence between the BIMJ and authors will, as far as possible via should be conducted via email quoting the reference number.

Conditions

Submission of an article for consideration for publication implies the transfer of the copyright from the authors to the BIMJ upon acceptance. The final decision of acceptance rests with the Editor-in-Chief. All accepted papers become the permanent property of the BIMJ and may not be published elsewhere without written permission from the BIMJ.

Ethics

Ethical considerations will be taken into account in the assessment of papers that have experimental investigations of human or animal subjects. Authors should state clearly in the Materials and Methods section of the manuscript that institutional review board has approved the project. Those investigators without such review boards should ensure that the principles outlined in the Declaration of Helsinki have been followed.

Manuscript categories

Original articles

These include controlled trials, interventional studies, studies of screening and diagnostic tests, outcome studies, cost-effectiveness analyses, and large-scale epidemiological studies. Manuscript should include the following; introduction, materials and methods, results and conclusion. The objective should be stated clearly in the introduction. The text should not exceed 2500 words and references not more than 30.

Review articles

These are, in general, invited papers, but unsolicited reviews, if of good quality, may be considered. Reviews are systematic critical assessments of

literature and data sources pertaining to clinical topics, emphasising factors such as cause, diagnosis, prognosis, therapy, or prevention. Reviews should be made relevant to our local setting and preferably supported by local data. The text should not exceed 3000 words and references not more than 40.

Special Reports

This section usually consist of invited reports that have significant impact on healthcare practice and usually cover disease outbreaks, management guidelines or policy statement paper.

Audits

Audits of relevant topics generally follow the same format as original article and the text should not exceed 1,500 words and references not more than 20.

Case reports

Case reports should highlight interesting rare cases or provide good learning points. The text should not exceed 1000 words; the number of tables, figures, or both should not be more than two, and references should not be more than 15.

Education section

This section includes papers (i.e. how to interpret ECG or chest radiography) with particular aim of broadening knowledge or serve as revision materials. Papers will usually be invited but well written paper on relevant topics may be accepted. The text should not exceed 1500 words and should include not more than 15 figures illustration and references

three relevant references should be included. Only images of high quality (at least 300dpi) will be acceptable.

Technical innovations

This section include papers looking at novel or new techniques that have been developed or introduced to the local setting. The text should not exceed 1000 words and should include not more than 10 figures illustration and references should not be more than 10.

Letters to the Editor

Letters discussing a recent article published in the BIMJ are welcome and should be sent to the Editorial Office by e-mail. The text should not exceed 250 words; have no more than one figure or table, and five references.

Criteria for manuscripts

Manuscripts submitted to the BIMJ should meet the following criteria: the content is original; the writing is clear; the study methods are appropriate; the data are valid; the conclusions are reasonable and supported by the data; the information is important; and the topic has general medical interest. Manuscripts will be accepted only if both their contents and style meet the standards required by the BIMJ.

Authorship information

Designate one corresponding author and provide a complete address, telephone and fax numbers, and e-mail address. The number of authors of each paper should not be more than twelve; a greater number requires justification. Authors may add a publishable footnote explaining order of authorship.

Group authorship

If authorship is attributed to a group (either solely or in addition to one or more individual authors), all members of the group must meet the full criteria and requirements for authorship described in the following paragraphs. One or more authors may take responsibility 'for' a group, in which case the other group members are not authors, but may be listed in an acknowledgement.

Authorship requirement

DISCLAIMER

All articles published, including editorials and letters, represent the opinion of the contributors and do not reflect the official view or policy of the Clinical Research Unit, the Ministry of Health or the institutions with which the contributors are affiliated to unless this is clearly stated. The appearance of advertisement does not necessarily constitute endorsement by the Clinical Research Unit or Ministry of Health, Brunei Darussalam. Furthermore, the publisher cannot accept responsibility for the correctness or accuracy of the advertisers' text and/or claim or any opinion expressed.

sign, and the analysis and interpretation of the data (where applicable); to have made substantial contributions to the writing or revision of the manuscript; and to have reviewed the final version of the submitted manuscript and approved it for publication. Authors will be asked to certify that their contribution represents valid work and that neither the manuscript nor one with substantially similar content under their authorship has been published or is being considered for publication elsewhere, except as described in an attachment. If requested, authors shall provide the data on which the manuscript is based for examination by the editors or their assignees.

Financial disclosure or conflict of interest

Any affiliation with or involvement in any organisation or entity with a direct financial interest in the subject matter or materials discussed in the manuscript should be disclosed in an attachment. Any financial or material support should be identified in the manuscript.

Copyright transfer

In consideration of the action of the BIMJ in reviewing and editing a submission, the author/s will transfer, assign, or otherwise convey all copyright ownership to the Clinical Research Unit, RIPAS Hospital, Ministry of Health in the event that such work is published by the BIMJ.

Acknowledgements

Only persons who have made substantial contributions but who do not fulfill the authorship criteria should be acknowledged.

Accepted manuscripts

Authors will be informed of acceptances and accepted manuscripts will be sent for copyediting. During copyediting, there may be some changes made to accommodate the style of journal format. Attempts will be made to ensure that the overall meaning of the texts are not altered. Authors will be informed by email of the estimated time of publication. Authors may be requested to provide raw data, especially those presented in graph such as bar charts or figures so that presentations can be constructed following the format and style of the journal. Proofs will be sent to authors to check for any mistakes made

KIDNEY HEALTH FOR ALL: BRIDGING THE GAP IN KIDNEY HEALTH EDUCATION AND LITERACY.

Robyn G. LANGHAM¹, Kamyar KALANTAR-ZADEH², Ann BONNER³, Alessandro BALDUC-
CI⁴, Li-Li HSIAO⁵, Latha A. KUMARASWAMI⁶, Paul LAFFIN⁷, Vassilios LIAKOPOULOS⁸,
Gamal SAADI⁹, Ekamol TANTISATTAMO², Ifeoma ULASI¹⁰ and Siu-Fai LUI¹¹ for the
World Kidney Day Joint Steering Committee¹².

¹St. Vincent's Hospital, Department of Medicine, University of Melbourne, Melbourne, Victoria, Australia.

²Division of Nephrology, Hypertension and Kidney Transplantation, Department of Medicine, University of California Irvine School of Medicine, Orange, California, USA.

³School of Nursing and Midwifery, Griffith University, Southport, Queensland, Australia.

⁴Italian Kidney Foundation, Rome, Italy.

⁵Brigham and Women's Hospital, Renal Division, Department of Medicine, Boston, Massachusetts, USA.

⁶Tamilnad Kidney Research (TANKER) Foundation, The International Federation of Kidney Foundations - World Kidney Alliance (IFKF - WKA), Chennai, India.

⁷International Society of Nephrology, Brussels, Belgium.

⁸Division of Nephrology and Hypertension, 1st Department of Internal Medicine, AHEPA Hospital, Aristotle University of Thessaloniki, Thessaloniki, Greece.

⁹Nephrology Unit, Department of Internal Medicine, Faculty of Medicine, Cairo University, Giza, Egypt.

¹⁰Renal Unit, Department of Medicine, College of Medicine, University of Nigeria, Ituku-Ozalla. Enugu, Nigeria.

¹¹International Federation of Kidney Foundations – World Kidney Alliance, The Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong, China.

¹²The World Kidney Day Joint Steering Committee is listed in the Appendix.

This article is being published in *Kidney International* and reprinted concurrently in several journals. The articles cover identical concepts and wording, but vary in minor stylistic and spelling changes, detail, and length of manuscript in keeping with each journal's style. Any of these versions may be used in citing this article.

Note that [all authors contributed equally to the conception, preparation, and editing of the manuscript.](#)

Copyright © 2022, World Kidney Day Steering Committee. Published by Elsevier Inc., on behalf of the International Society of Nephrology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Corresponding author: Robyn G. Langham, MBBS, PhD, University of Melbourne, Melbourne, Victoria, Australia. Tel: +61 414851892. E-mail: rlangham@unimelb.edu.au

Reprints: Ekamol Tantisattamo, Division of Nephrology, Hypertension and Kidney Transplantation, Department of Medicine, University of California Irvine School of Medicine, 101 The City Drive South, Orange, California 92868, USA. Tel: +1-714-456-5142. Fax: +1-714-456-6034. E-mail: etantisa@hs.uci.edu

ABSTRACT

The high burden of kidney disease, global disparities in kidney care, and poor outcomes of kidney failure bring a concomitant growing burden to persons affected, their families, and carers, and the community at large. Health literacy is the degree to which persons and organizations have or equitably enable individuals to have the ability to find, understand, and use information and services to make informed health-related decisions and actions for themselves and others. Rather than viewing health literacy as a patient deficit, improving health literacy largely rests with health care providers communicating and educating effectively in codesigned partnership with those with kidney disease. For kidney policy makers, health literacy provides the imperative to shift organizations to a culture that places the person at the center of health care. The growing capability of and access to technology provides new opportunities to enhance education and awareness of kidney disease for all stakeholders. Advances in telecommunication, including social media platforms, can be leveraged to enhance persons' and providers' education; The World Kidney Day declares 2022 as the year of "Kidney Health for All" to promote global teamwork in advancing strategies in bridging the gap in kidney health education and literacy. Kidney organizations should work toward shifting the patient-deficit health literacy narrative to that of being the responsibility of health care providers and health policy makers. By engaging in and supporting kidney health-centered policy making, community health planning, and health literacy approaches for all, the kidney communities strive to prevent kidney diseases and enable living well with kidney disease.

KEYWORDS: Educational gap, Empowerment, Health literacy, Health policy, Information technology, Kidney health, Partnership, Prevention, Social media.

INTRODUCTION

Given the high burden of kidney disease and global disparities related to kidney care, in carrying forward our mission of advocating *Kidney Health for All*, the challenging issue of bridging the well-identified gap in the global understanding of kidney disease and its health literacy is the theme for World Kidney Day (WKD) 2022. Health literacy is defined as the degree to which persons and organizations have—or equitably enable individuals to have—the ability to find, understand, and use information and services to inform health-related decisions and actions for themselves and others.¹ Not only is there is growing recognition of the role that health literacy has in determining outcomes for persons affected by kidney disease and the community in general, but there is an emergent imperative for policy makers worldwide to be informed and cognizant of opportunities and real measurable outcomes that can be achieved through kidney-specific preventative strategies.

THE GLOBAL COMMUNITY OF PEOPLE WITH KIDNEY DISEASE

Most people are not aware of what kidneys are for or even where their kidneys are. For those afflicted by disease and the subsequent effects on overall health, an effective health care provider communication is required to support individuals to be able to understand what to do, to make decisions, and to take action. Health literacy involves more than functional abilities of an individual; it is also the cognitive and social skills needed to gain access to, understand, and use information to manage health condition.² It is also contextual³ in that as health needs change, so too does the level of understanding and ability to problem solve alter. Health literacy is, therefore, an interaction between individuals, health care providers, and health policy makers.⁴ This why the imperatives around health literacy are now recognized as indicators for the quality of local and national health care systems and health care professionals within

it.⁵ For chronic kidney disease (CKD), as the disease progresses alongside other health changes and increasing treatment complexities, it becomes more difficult for individuals to manage.⁶ Promoted in health policy for around a decade involving care partnerships between health-centered policy, community health planning, and health literacy,⁷ current approaches need to be shifted forward (Table I).

Assessing health literacy necessitates the use of appropriate multidimensional patient-reported measures, such as the World Health Organization–recommended Health Literacy Questionnaire (available in over 30 languages) rather than tools measuring only functional health literacy (e.g., Rapid Estimate of Adult Literacy in Medicine or Short Test of Functional Health Literacy in Adults).⁸ It is therefore not surprising that studies of low health literacy (LHL) abilities in people

Table I: Summary characteristic of kidney health promotion, involving kidney health–centered policy, community kidney health planning, and kidney health literacy, and proposed future direction.

Kidney health promotion	Definition	Stakeholders	Current status	Limitations/ challenges	Suggested solutions/future research
Kidney health–centered policy	<ul style="list-style-type: none"> • Incorporate kidney health into policy decision making • Prioritize policies with primary prevention for CKD 	<ul style="list-style-type: none"> • Governance • Policy makers • Insurance agencies 	<ul style="list-style-type: none"> • Policy emphasizing treatment for CKD and kidney failure rather than kidney health prevention 	<ul style="list-style-type: none"> • Economic-driven situation challenging CKD risk factor minimization (e.g., food policy) 	<ul style="list-style-type: none"> • Promote implementation of public health program for primary CKD prevention • Promote sustainable treatment for CKD and dialysis • - Increase kidney transplant awareness • - Enhance visibility and encourage brother-sister nephrology and transplant program in LMIC • Support research funding from government • - Health care cost-effectiveness for caring for CKD • - Kidney failure, including maintenance dialysis and transplant • - Promote surveillance programs for kidney diseases and their risk factors
Community kidney health planning	<ul style="list-style-type: none"> • Building up preventive strategies to promote healthy communities and primary health care facilities 	<ul style="list-style-type: none"> • Community leadership • Kidney patient advocacy 	<ul style="list-style-type: none"> • Belief in community leaders in LMIC 	<ul style="list-style-type: none"> • Education and understanding kidney health promotion of community leadership and people 	<ul style="list-style-type: none"> • Improve role model of community • Enhance kidney support networks
Kidney health literacy	<ul style="list-style-type: none"> • Receive knowledge, skills, and information to be healthy 	<ul style="list-style-type: none"> • People with CKD • Care partners • Health care providers 	<ul style="list-style-type: none"> • Lack of awareness of CKD and risk factors • Care partner burden and burnout • Inadequate health care workers • High patients-to-health care workers ratio, especially in rural areas 	<ul style="list-style-type: none"> • Inadequate policy direction • Ineffective health care providers’ communication skills 	<ul style="list-style-type: none"> • Organizational paradigm shift toward health literacy • Improving communication between health care providers with patients and care partners • Using teach-back methods for consumer education • Adapting technologies for appropriate health literacy and socio-cultural environments • Family engagement in the patient care • Incentive for community health care providers in rural areas

CKD: Chronic Kidney Disease; LMIC: Low-to middle-Income Country.

with CKD have been demonstrated to be associated with poor CKD knowledge, self-management behaviors, and health-related quality of life and in those with greater comorbidity severity.⁷ Unfortunately, most CKD studies have measured only functional health literacy, so the evidence that LHL results in poorer outcomes, particularly that it increases health care utilization and mortality,⁹ and reduces access to transplantation,¹⁰ is weak.

Recently, health literacy is now considered to be an important bridge between lower socioeconomic status and other social determinants of health.⁴ Indeed, this is not a feature that can be measured by the gross domestic product of a country, as the effects of LHL on the extent of CKD in the community are experienced globally regardless of country income status. The lack of awareness of risk factors of kidney disease, even in those with high health literacy abilities, is testament to the difficulties in understanding this disease, and why the United States, for instance, recommends that a universal precautions approach toward health literacy is undertaken.¹¹

So, what does the perfect health literacy program look like for people with CKD? In several high-income countries, there are national health literacy action plans with the emphasis shifted to policy directives, organizational culture, and health care providers. In Australia, for instance, a compulsory health literacy accreditation standard makes the health care organization responsible for ensuring providers are cognizant of individual health literacy abilities.¹² Although many high-income countries, health care organizations, nongovernmental organizations, and jurisdictions are providing an array of consumer-facing web-based programs that provide detailed information and self-care training opportunities, most are largely designed for individual/family use that are unlikely to miti-

gate LHL. There is, however, substantial evidence that interventions improving health care provider communication are more likely to improve understanding of health problems and abilities to adhere to complex treatment regimens.¹³

Access to information that is authentic and tailored specifically to the needs of the individual and the community is the aim. The challenge is recognized acutely in more remote and low- to middle-income countries of the world, specifically the importance of culturally appropriate knowledge provision. The principals of improving health literacy are the same, but understanding how to proceed, and putting consumers in charge, with a codesign approach, is critical and may result in a different outcome in more remote parts of the world. This principal especially applies to communities that are smaller, with less access to electronic communication and health care services, where the level of health literacy is shared across the community and where what affects the individual also affects all the community. Decision support systems are different, led by elders, and in turn educational resources are best aimed at improving knowledge of the whole community.

A systematic review of the evaluation of interventions and strategies shows this area of research is still at an early stage,¹⁴ with no studies unravelling the link between LHL and poor CKD outcomes. The best evidence is in supporting targeted programs on improving communication capabilities of health care professionals as central. One prime example is Teach-back, a cyclical, simple, low-cost education intervention, shows promise for improving communication, knowledge, and self-management in the CKD populations in low- or high-income countries.¹⁵ Furthermore, the *consumer*-led voice has articulated research priorities that align closely with principals felt to be important to success of education: building new education

THE NETWORKED COMMUNITY OF KIDNEY HEALTH CARE WORKERS

Nonphysician health care workers, including nurses and advanced practice providers (physician assistants and nurse practitioners) as well as dietitians, pharmacists, social workers, technicians, physical therapists, and other allied health professionals, often spend more time with persons with kidney disease, compared with nephrologists and other physician specialists. In an ambulatory care setting at an appointment, in the emergency department, or in the inpatient setting, these health care professionals often see and relate to the patient first, last, and in between, given that physician encounters are often short and focused. Hence, the nonphysician health care workers have many opportunities to discuss kidney disease-related topics with the individuals and their care partners and to empower them.^{17,18} For instance, medical assis-

tants can help identify those with or at risk of developing CKD and can initiate educating them and their family members about the role of diet and lifestyle modification for primary, secondary, and tertiary prevention of them and their family members about the role of diet and lifestyle modification for primary, secondary, and tertiary prevention of CKD while waiting to see the physician.¹⁹ Some health care workers provide networking and support for kidney patient advocacy groups and kidney support networks, which have been initiated or expanded via social media platforms (Figure 1).^{20,21} Studies examining the efficacy of social media in kidney care and advocacy are on the way.^{22,23}

Like physicians, many activities of nonphysician health care workers have been increasingly affected by the rise of electronic health recording and growing access to internet-based resources, including social media, that offer educational materials related to




Figure 1: Schematic representation of consumer and health care professionals' collaborative advocacy using social media platforms with the goal of *Kidney Health for All*.

Table II: Social media that are more frequently used for kidney education and advocacy.

Social media	Strength	Limitations	Additional comments
Facebook	⇒ Frequently used social media platform by many kidney patients and patient groups	⇒ Widely used for entertaining purposes, which can dilute its professional utility	⇒ User-friendly platform for kidney advocacy, enabling wide ranges of outreach goals
Instagram	⇒ Photo-predominating platform	⇒ Not frequently used by health care professionals	⇒ Picture friendly, potentially effective for illustrative educational purposes
Twitter	⇒ Often used by physician specialists and scientists, including nephrologists	⇒ Less frequently used by patients and care partners	⇒ Increasing popularity among physician and specialty circles
LinkedIn	⇒ More often used by professionals, including in industry	⇒ Originally designed for employment and job-seeking networking	⇒ Mostly effective to reach out to industry and managerial professionals
YouTube	⇒ Video-predominating platform	⇒ Less effective with non-video-based formats	⇒ Wide ranges of outreach and educational targets
WeChat	⇒ Widely used in mainland China	⇒ Access is often limited to those living in China or its diaspora	⇒ Effective platform to reach out to patients and health care professionals in China
Pinterest	⇒ Picture-based, often used by dietitians	⇒ Currently limited use by some health care workers	⇒ Useful for dietary and lifestyle education

Other popular social media at the time of this publication include, but not limited to, Tik Tok, Snapchat, Reddit, Tumblr, Telegram, Quora, and many others that are currently only occasionally used in kidney advocacy activities. Mobile and social media messaging apps include, but not limited to, WhatsApp, Zoom, Facebook Messengers, Skype Teams, and Slack. Note that platforms that are more often used as internet-based messaging are not included.

kidney health, including kidney-preserving therapies with traditional and emerging interventions.²⁴ These resources can be used for both self-education and for networking and advocacy on kidney disease awareness and learning. Increasingly, more health care professionals are engaged in some types of social media-based activities, as shown in Table II. At the time of this writing, the leading social media used by many—but not all—kidney health care workers include Facebook, Instagram, Twitter, LinkedIn, and YouTube. In some regions of the world, certain social media are more frequently used than others given unique cultural or access constellations (e.g., WeChat is a platform often used by health care workers and patient groups in China). Some health care professionals, such as managers and those in leadership and advocacy organization positions, may choose to embark on social media to engage those with CKD and their care partners or other health care professionals in alliance building and marketing. To that end, effective communication strategies and outreach skills specific to responsible use of social media can provide

clear advantages given that these skills and strategies are different and may need modification in those with LHL. It is imperative to ensure the needed knowledge and training for accountable approach to social media is provided to health care providers, so that these outreach strategies are utilized with the needed awareness of their unique strengths and pitfalls, as follows²⁵:

- (i) Consumers' and care partners' confidentiality may not be breached upon posting anything on social media, including indirect referencing to a specific individual or a particular description of a condition unique to a specific person (e.g., upon soliciting for transplant kidney donors on social media).^{26,27}
- (ii) Confidential information about clinics, hospitals, dialysis centers, or similar health care and advocacy entities may not be disclosed on social media without ensuring that the needed processes, including collecting authorizations to disclose, are undertaken.
- (iii) Health care workers' job security and ca-

reers should remain protected with thorough review of the content of the messages and illustrations/videos before online posting.

- (iv) Careless and disrespectful language and emotional tones are often counterproductive and may not be justified under the context of freedom of speech.

THE GLOBAL KIDNEY COMMUNITY OF POLICY AND ADVOCACY

Policy and advocacy are well-recognized tools that, if properly deployed, can bring about change and paradigm shift at jurisdictional level. The essence of advocating for policy change to better address kidney disease is, in itself, an exercise in improving health literacy of the policy makers. Policy development, at its core, is a key stakeholder or stakeholder group (e.g., the kidney community, who believes that a problem exists that should be tackled through governmental action). There is an increasing recognition of the importance of formulating succinct, meaningful, and authentic information, akin to improving health literacy, to present to government for action.

Robust and efficacious policy is always underpinned by succinct and applicable information; however, the development and communication of this message, designed to bridge the gap in knowledge of relevant jurisdictions, is only part of the process of policy development. An awareness of the process is important to clinicians who are aiming to advocate for effective change in prevention or improvement of outcomes in the CKD community.

Public policies, the plans for future action accepted by governments, are articulated through a political process in response stakeholder observation, usually written as a directive, law, regulation, procedure, or circular. Policies are purpose fit and targeted to

defined goals and specific societal problems and are usually a chain of actions effected to solve those societal problems.²⁸ Policies are an important output of political systems. Policy development can be formal, passing through rigorous lengthy processes before adoption (such as regulations), or it can be less formal and quickly adopted (such as circulars). As already mentioned, the governmental action envisaged by the key stakeholders as solution to a problem is at its core. The process enables stakeholders to air their views and bring their concerns to the fore. Authentic information that is meaningful to the government is critical. The policy development process can be stratified into 5 stages (i.e., the policy cycle), as depicted by Andersen (1994)²⁹ and adapted and modified by other authors³⁰ (Figure 2). The policy cycle constitutes an expedient framework for evaluating the key components of the process.

Subsequently, the policy moves on to implementation phase. This phase may require subsidiary policy development and adoption of new regulations or budgets (implementation). Policy evaluation is integral to the policy processes and applies evaluation principles and methods to assess the content, implementation, or impact of a policy. Evaluation facilitates understanding and appreciation of the worth and merit of a policy as well as the need for its improvement. More important, of the 5 principles of advocacy that underline policy making,³¹ the most important for clinicians engaged in this space is that of commitment, persistence, and patience. Advocacy takes time to yield the desired results.

The Advocacy Planning Framework, developed by Young and Quinn in 2002,³⁰ consists of overlapping circles representing 3 sets of concepts (way into the process, the messenger, and message and activities) that are key to planning any advocacy campaign:

- (i) "Way into the process": discusses the best




Figure 2: Policy cycle involving 5 stages of policy development. CKD, chronic kidney disease; RRT, kidney replacement therapy; LGA, local government area.

approaches to translate ideas into the target policy debate and identify the appropriate audience to target.

- (ii) Messenger: talks about the image maker or face of the campaign and other support paraphernalia that are needed.
- (iii) Message and activities: describe what can be said to the key target audiences that is engaging and convincing. And how best it can be communicated through appropriate communication tools.

Advocacy is defined as "an effort or campaign with a structured and sequenced plan of action which starts, directs, or prevents a specific policy change."³¹ The goal being to influence decision makers through communicating directly with them or getting their commitment through secondary audiences (advisers, the media, or the public) to

the end that the decision maker understands, is convinced, takes ownership of the ideas, and finally has the compulsion to act.³¹ As with improving health literacy, it is the communication of ideas to policy makers for adoption and implementation as policy that is key. There is much to be done with bridging this gap in understanding of the magnitude of community burden that results from CKD. Without good communication, many good ideas and solutions do not reach communities and countries where they are needed. Again, aligned with the principles of developing resources for health literacy, the approach also needs to be nuanced according to the local need, aiming to have the many good ideas and solutions be communicated to communities and countries where they are needed.

Advocacy requires galvanizing momen-

tum and support for the proposed policy or recommendation. The process is understandably slow as it involves discussions and negotiations for paradigms, attitudes, and positions to shift. In contemplating advocacy activities, multiple factors must be considered, interestingly not too dissimilar to that of building health literacy resources: What obstructions are disrupting the policy-making process from making progress? What resources are available to enable the process to succeed? Is the policy objective achievable considering all variables? Is the identified problem already being considered by the policy makers (government or multinational organizations)? Any interest or momentum generated around it? Understandably, if there is some level of interest and if government already has its spotlight on the issue, it is likely to succeed.

Approaches to choose from include the following^{31,32}:

- Advising (researchers are commissioned to produce new evidence-based proposals to assist the organization in decision making).
- Activism: involves petitions, public demonstrations, posters, fliers, and leaflet dissemination, often used by organizations to promote a certain value set.
- Media campaign: having public pressure on decision makers helps in achieving results.
- Lobbying: entails face-to-face meetings with decision makers; often used by business organizations to achieve their purpose.

Here lies the importance of effective and successful advocacy to stakeholders, including policy makers, health care professionals, communities, and key change makers in society. The WKD, since inception, has aimed at playing this role. WKD has gained people's trust by delivering relevant and accurate

messaging and supporting leaders in local engagement, and it is celebrated by kidney care professionals, celebrities, those with the disease, and their care givers all over the world. To achieve the goal, an implementation framework of success in a sustainable way includes creativity, collaboration, and communication.

The ongoing challenge for the International Society of Nephrology and International Federation of Kidney Foundations–World Kidney Alliance, through the Joint Steering Committee of WKD, is to operationalize how to collate key insights from research and analysis to effectively feed the policy-making process at the local, national, and international levels, to inform or guide decision making (i.e., increasing engagement of governments and organizations, like World Health Organization, United Nations, and regional organizations, especially in low-resource settings). There is a clear need for ongoing renewal of strategies to increase efforts at closing gap in kidney health literacy, empowering those affected with kidney disease and their families, giving them a voice to be heard, and engaging with the civil society. This year, the Joint Steering Committee of WKD declares “*Kidney Health for All*” as the theme of the 2022 WKD to emphasize and extend collaborative efforts among people with kidney disease, their care partners, health care providers, and all involving stakeholders for elevating education and awareness on kidney health and saving lives with this disease.

CONCLUSIONS

In bridging the gap of knowledge to improve outcomes for those with kidney disease on a global basis, an in-depth understanding of the needs of the community is required. The same can be said for policy development, understanding the processes in place for engagement of governments worldwide, all un-

derstanding the processes in place for engagement of governments worldwide, all underpinned by the important principal of codesign of resources and policy that meets the needs of the community for which it is intended.

For World Kidney Day 2022, kidney organizations, including the International Society of Nephrology and International Federation of Kidney Foundations–World Kidney Alliance, have a responsibility to immediately work toward shifting the patient-deficit health literacy narrative to that of being the responsibility of clinicians and health policy makers. LHL occurs in all countries regardless of income status; hence, simple, low-cost strategies are likely to be effective. Communication, universal precautions, and teach back can be implemented by all members of the kidney health care team. Through this vision, kidney organizations will lead the shift to improved patient-centered care, support for care partners, health outcomes, and the global societal burden of kidney health care.

ORCID id

0000-0002-2735-0161 Robyn G. Langham
 0000-0002-8666-0725 Kamyar Kalantar-Zadeh
[0000-0001-9920-6743](#) Ann Bonner
 0000-0002-7432-1494 Alessandro Balducci
 0000-0003-0830-3391 Li-Li Hsiao
 0000-0002-9230-2473 Paul Laffin
 0000-0002-7564-2724 Vassilios Liakopoulos
 0000-0002-6280-1685 Gamal Saadi
 0000-0003-0883-6892 Ekamol Tantisattamo
 0000-0001-7783-3025 Ifeoma Ulasi
 0000-0003-0800-8982 Siu-Fai Lui

CONFLICT OF INTEREST DISCLOSURE

KK-Z reports honoraria from Abbott, Abbvie, ACI Clinical, Akebia, Alexion, Amgen, Ardelyx, AstraZeneca, Aveo, BBraun, Cara Therapeu-

tics, Chugai, Cytokinetics, Daiichi, DaVita, Fresenius, Genentech, Haymarket Media, Hospira, Kabi, Keryx, Kissei, Novartis, Pfizer, Regulus, Relypsa, Resverlogix, Dr Schaer, Sandoz, Sanofi, Shire, Vifor, UpToDate, and ZS-Pharma.

VL reports nonfinancial support from Genesis Pharma.

GS reports personal fees from Multicare, Novartis, Sandoz, and AstraZeneca.

ET reports nonfinancial support from Natera.

All the other authors declared no competing interests.

APPENDIX

The World Kidney Day Joint Steering Committee includes coauthors Robyn G. Langham, Kamyar Kalantar-Zadeh, Alessandro Balducci, Li-Li Hsiao, Latha Kumaraswami, Paul Laffin, Vassilios Liakopoulos, Gamal Saadi, Ifeoma Ulasi, and Siu-Fai Lui.

REFERENCES

- Centers for Disease Control and Prevention. [Healthy People 2030: What Is Health Literacy?](#) [Accessed January 16, 2022].
- Nutbeam D. The evolving concept of health literacy. *Soc Sci Med.* 2008;67(12):2072-8. doi:10.1016/j.socscimed.2008.09.050
- Lloyd A, Bonner A, Dawson-Rose C. [The health information practices of people living with chronic health conditions: Implications for health literacy.](#) *J Librarianship Information Science.* 2014;46:207-2016. [Accessed January 16, 2022].
- Sorensen K, Van den Broucke S, Fullam J, et al. [Health literacy and public health: a systematic review and integration of definitions and models.](#) *BMC Public Health.* 2012;12:80. doi:10.1186/1471-2458-12-80. [Accessed January 16, 2022].
- Nutbeam D, Lloyd JE. [Understanding and Responding to Health Literacy as a Social Determinant of Health.](#) *Annu Rev Public Health.* 2021;42:159-173. doi:10.1146/annurev-publhealth-090419-102529. [Accessed January 16, 2022].

- 6: Mathias-Shah J, Ramsbotham J, Seib C, Muir R, Bonner A. A scoping review of the role of health literacy in chronic kidney disease self-management. *J Ren Care.* 2021;47:221-233.
- 7: Dinh HTT, Nguyen NT, Bonner A. [Healthcare systems and professionals are key to improving health literacy in chronic kidney disease.](#) *J Ren Care.* 2021;doi:10.1111/jorc.12395. [Accessed January 16, 2022].
- 8: Dobson S, Good S, Osborne R. [Health literacy toolkit for low and middle-income countries: A series of information sheets to empower communities and strengthen health systems \[Internet\].](#) *New Delhi: World Health Organization.* 2015. [Accessed January 16, 2022].
- 9: Taylor DM, Fraser S, Dudley C, et al. [Health literacy and patient outcomes in chronic kidney disease: a systematic review.](#) *Nephrol Dial Transplant.* 2018;33(9):1545-1558. doi:10.1093/ndt/gfx293. [Accessed January 16, 2022].
- 10: Taylor DM, Bradley JA, Bradley C, et al. [Limited health literacy is associated with reduced access to kidney transplantation.](#) *Kidney Int.* 2019;95(5):1244-1252. doi:10.1016/j.kint.2018.12.021. [Accessed January 16, 2022].
- 11: Brega AG, Barnard J, Mabachi NM, et al. [AHRQ Health Literacy Universal Precautions Toolkit, Second Edition.](#) (Prepared by Colorado Health Outcomes Program, University of Colorado Anschutz Medical Campus under Contract No. HHS290200710008, TO#10.) AHRQ Publication No. 15-0023-EF. Rockville, MD: Agency for Healthcare Research and Quality; January 2015. [Accessed January 16, 2022].
- 12: Australian Commission on Safety and Quality in Health Care. [Health literacy: Taking action to improve safety and quality.](#) Sydney: ACSQHC, 2014. [Accessed January 17, 2022].
- 13: Visscher BB, Steunenberg B, Heijmans M, et al. [Evidence on the effectiveness of health literacy interventions in the EU: a systematic review.](#) *BMC Public Health.* 2018;18(1):1414. doi:10.1186/s12889-018-6331-7. [Accessed January 16, 2022].#
- 14: Boonstra MD, Reijneveld SA, Foitzik EM, Westerhuis R, Navis G, de Winter AF. [How to tackle health literacy problems in chronic kidney disease patients? A systematic review to identify promising intervention targets and strategies.](#) *Nephrol Dial Transplant.* 2021;36:1207-1221. doi:10.1093/ndt/gfaa273. [Accessed January 16, 2022].
- 15: Nguyen NT, Douglas C, Bonner A. Effectiveness of self-management programme in people with chronic kidney disease: A pragmatic randomized controlled trial. *J Adv Nurs.* Mar 2019;75(3):652-664. doi:10.1111/jan.13924
- 16: Synnot A, Bragge P, Lowe D, et al. [Research priorities in health communication and participation: international survey of consumers and other stakeholders.](#) *BMJ Open.* 05 08 2018;8(5):e019481. doi:10.1136/bmjopen-2017-019481. [Accessed January 17, 2022].
- 17: Kalantar-Zadeh K, Kam-Tao Li P, Tantisattamo E, et al. [Living well with kidney disease by patient and care-partner empowerment: kidney health for everyone everywhere.](#) *Brunei Int Med J.* 2021;17:17-25. [Accessed January 17, 2022].
- 18: Jager KJ, Kovesdy C, Langham R, Rosenberg M, Jha V, Zoccali C. [A single number for advocacy and communication-worldwide more than 850 million individuals have kidney diseases.](#) *Kidney Int.* 2019;96(5):1048-1050. doi:10.1016/j.kint.2019.07.012. [Accessed January 17, 2022].
- 19: Li PK, Garcia-Garcia G, Lui SF, et al. [Kidney health for everyone everywhere-from prevention to detection and equitable access to care.](#) *Brunei Int Med J.* 2020;16:60-72. [Accessed January 17, 2022].
- 20: Gilford S. Patients helping patients: the Renal Support Network. *Nephrol Nurs J.* 2007;34(1):76.
- 21: Muhammad S, Allan M, Ali F, Bonacina M, Adams M. [The renal patient support group: supporting patients with chronic kidney disease through social media.](#) *J Ren Care.* 2014;40(3):216-8. doi:10.1111/jorc.12076. [Accessed January 17, 2022].
- 22: Li WY, Chiu FC, Zeng JK, et al. [Mobile Health App With Social Media to Support Self-Management for Patients With Chronic Kidney Disease: Prospective Randomized Controlled Study.](#) *J Med Internet Res.* 2020;22(12):e19452. doi:10.2196/19452. [Accessed January 17, 2022].
- 23: Pase C, Mathias AD, Garcia CD, Garcia Rodrigues C. [Using Social Media for the Promotion of Education and Consultation in Adolescents Who Have Undergone Kidney Transplant: Protocol for a Randomized Control Trial.](#) *JMIR Res Protoc.* 2018;7(1):e3. doi:10.2196/resprot.8065. [Accessed January 17, 2022].

- 24: Kalantar-Zadeh K, Jafar TH, Nitsch D, Neuen BL, Perkovic V. Chronic kidney disease. *Lancet*. 2021;398(10302):786-802. doi:10.1016/S0140-6736(21)00519-5
 - 25: Chen L, Sivaparthipan CB, Rajendiran S. Un-professional problems and potential healthcare risks in individuals' social media use. *Work*. 2021;68(3):945-953. doi:10.3233/WOR-203428
 - 26: Henderson ML, Herbst L, Love AD. Social Media and Kidney Transplant Donation in the United States: Clinical and Ethical Considerations When Seeking a Living Donor. *Am J Kidney Dis*. 2020;76(4):583-585. doi:10.1053/j.ajkd.2020.03.027
 - 27: #Henderson ML. [Social Media in the Identification of Living Kidney Donors: Platforms, Tools, and Strategies](#). *Curr Transplant Rep*. 2018;5(1):19-26. [Accessed January 17, 2022].
 - 28: Newton K, van Deth JW, eds. [Foundations of Comparative Politics Democracies of the Modern World](#). 2nd ed. Cambridge, UK: Cambridge University Press; 2010. [Accessed December 13, 2021].
 - 29: Anderson JE. *Public Policymaking: An Introduction*. 2nd ed. Boston, MA: Houghton Mifflin; 1994.
 - 30: Young E, Quinn L, eds. [Writing Effective Public Policy Papers: A Guide to Policy Advisers in Central and Eastern Europe](#). Budapest, Hungary: Open Society Institute; 2002. [Accessed December 13, 2021].
 - 31: Young E, Quinn L, eds. [Making Research Evidence Matter: A Guide to Policy Advocacy in Transition Countries](#). Budapest, Hungary: Open Society Foundations; 2012. [Accessed December 13, 2021].
 - 32: Start D, Hovland I. [Tools for Policy Impact: A Handbook for Researchers, Research and Policy in Development Programme](#). London, UK: Overseas Development Institute; 2004. [Accessed December 13, 2021].
-